


This article was downloaded by:[McMaster University Library]
On: 2 September 2007
Access Details: [subscription number 769426004]
Publisher: Taylor & Francis
Informa Ltd Registered in England and Wales Registered Number: 1072954
Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



Optimization Methods and Software

Publication details, including instructions for authors and subscription information:
<http://www.informaworld.com/smpp/title~content=t713645924>

The central path visits all the vertices of the Klee-Minty cube

Antoine Deza^a; Eissa Nematollahi^a; Reza Peyghami^a; Tamás Terlaky^a

^a Advanced Optimization Laboratory, Department of Computing and Software, McMaster University, Hamilton, Ontario, Canada

Online Publication Date: 01 October 2006

To cite this Article: Deza, Antoine, Nematollahi, Eissa, Peyghami, Reza and Terlaky, Tamás (2006) 'The central path visits all the vertices of the Klee-Minty cube', Optimization Methods and Software, 21:5, 851 - 865

To link to this article: DOI: 10.1080/10556780500407725

URL: <http://dx.doi.org/10.1080/10556780500407725>

PLEASE SCROLL DOWN FOR ARTICLE

Full terms and conditions of use: <http://www.informaworld.com/terms-and-conditions-of-access.pdf>

This article maybe used for research, teaching and private study purposes. Any substantial or systematic reproduction, re-distribution, re-selling, loan or sub-licensing, systematic supply or distribution in any form to anyone is expressly forbidden.

The publisher does not give any warranty express or implied or make any representation that the contents will be complete or accurate or up to date. The accuracy of any instructions, formulae and drug doses should be independently verified with primary sources. The publisher shall not be liable for any loss, actions, claims, proceedings, demand or costs or damages whatsoever or howsoever caused arising directly or indirectly in connection with or arising out of the use of this material.

© Taylor and Francis 2007

The central path visits all the vertices of the Klee–Minty cube

ANTOINE DEZA, EISSA NEMATOLLAHI, REZA PEYGHAMI and
TAMÁS TERLAKY*

Advanced Optimization Laboratory, Department of Computing and Software,
McMaster University, Hamilton, Ontario, Canada

(Received 23 June 2005, in final form 6 October 2005)

The Klee–Minty cube is a well-known worst-case example for which the simplex method takes an exponential number of iterations as the algorithm visits all the 2^n vertices of the n -dimensional cube. While such behaviour is excluded by polynomial interior point methods, we show that, by adding an exponential number of redundant inequalities, the central path can be bent along the edges of the Klee–Minty cube. More precisely, for an arbitrarily small δ , the central path takes $2^n - 2$ turns as it passes through the δ -neighbourhood of all the vertices of the Klee–Minty cube in the same order as the simplex method does.

Keywords: Linear programming; Central path; Klee–Minty cube

1. Introduction

While the *simplex method*, introduced by Dantzig [1] works very well in practice for linear optimisation problems, Klee and Minty [2] gave an example in 1972 for which the simplex method takes an exponential number of iterations. More precisely, they considered a maximisation problem over an n -dimensional squashed cube and proved that a variant of the simplex method visits all of its 2^n vertices, that is, the time complexity is not polynomial for the worst case, as $2^n - 1$ iterations are necessary for this n -dimensional linear optimisation problem. The pivot rule used in the Klee–Minty example was the most negative reduced cost, but variants of the Klee–Minty n -cube showing an exponential running time exist for most pivot rules, see [3] and the references therein. The Klee–Minty worst-case example partially stimulated the search for a polynomial algorithm and, in 1979, Khachiyan's [4] *ellipsoid method* proved that linear programming is indeed polynomially solvable. In 1984, Karmarkar [5] proposed a more efficient polynomial algorithm that sparked the research on polynomial *interior point methods*. In short, while the simplex method goes along the edges of the polyhedron corresponding to the feasible region, interior point methods pass through the interior of this polyhedron. Starting at

*Corresponding author. Email: terlaky@mcmaster.ca

the *analytic centre*, most interior point methods follow the so-called *central path* and converge to the analytic centre of the optimal face, see for example, [6–8,11,12].

In this paper, following the Klee–Minty approach, we show that, by carefully adding an exponential number of redundant constraints to the Klee–Minty n -cube, the central path can be bent along its edges. In other words, we give an example where, for an arbitrarily small δ , starting from the δ -neighbourhood of a vertex adjacent to the optimal solution, the central path takes $2^n - 2$ turns as, before converging to the optimal solution, it passes through the δ -neighbourhood of all the vertices of the Klee–Minty cube in the same order as the simplex method does.

Before stating the main result in section 2 and giving its proof in section 3, we illustrate the bending of the central path in the two and three dimensional cases. Figures 1 and 2 show the trajectory of the central path starting from the highest vertex and converging to the origin after visiting each vertex of the Klee–Minty cube. The redundant constraints correspond to hyperplanes parallel to the facets of the cube containing the origin. More precisely, in dimension 2, the redundant inequality $16 + x_1 \geq 0$ is added 15,360 times and the redundant inequality $16 + x_2 - x_1/4 \geq 0$ is added 40,960 times. Starting from the highest vertex and with $\delta = 0.1$, the central path visits the δ -neighbourhood of each vertex of the Klee–Minty cube in the same order as the simplex algorithm does before converging to the optimal solution, that is, the origin. In dimension 3, the redundant inequality $48 + x_1 \geq 0$ (resp. $48 + x_2 - x_1/4 \geq 0$ and $48 + x_3 - x_2/4 \geq 0$) is added 161,280 (resp. 552,960, and 1,474,560 times).




Figure 1. Central path nearing all the vertices of the Klee–Minty 2-cube.




Figure 2. Central path nearing all the vertices of the Klee–Minty 3-cube.

2. Notations and the main result

We consider the following Klee–Minty variant where ε is a small positive factor by which the unit cube $[0, 1]^n$ is squashed.

$$\begin{aligned} \min \quad & x_n \\ \text{subject to} \quad & 0 \leq x_1 \leq 1 \\ & \varepsilon x_{k-1} \leq x_k \leq 1 - \varepsilon x_{k-1} \quad \text{for } k = 2, \dots, n. \end{aligned}$$

We denote this linear optimisation problem by KM. This minimisation problem has $2n$ constraints, n variables and the feasible region is an n -dimensional cube denoted by \mathcal{C} . Some variants of the simplex method take $2^n - 1$ iterations to solve KM as they visit all the vertices ordered by the decreasing value of the last coordinate x_n starting from $v^{(n)} = (0, \dots, 0, 1)$ until the optimal value $x_n^* = 0$ is reached at the origin v^θ . If an interior point method is used to solve KM, the central path starts from the analytic centre χ of \mathcal{C} and converges to the origin, as shown in figure 3.

While adding a set h of redundant inequalities does not change the feasible region of KM, the analytic centre χ^h and the central path are affected by the addition of redundant constraints. We consider redundant inequalities induced by hyperplanes parallel to the n facets of \mathcal{C} containing the origin. To ease the analysis, we consider that all redundant hyperplanes are put at the same distance d to the corresponding parallel facet of \mathcal{C} . The constraint parallel to $H_1: x_1 = 0$ is added h_1 times and the constraint parallel to $H_k: x_k = \varepsilon x_{k-1}$ is added h_k times for $k = 2, \dots, n$. By abuse of notation, the set h is denoted by the integer vector $h = (h_1, \dots, h_n)$. With these notations, the redundant linear optimisation problem KM^h is defined by

$$\begin{aligned} \min \quad & x_n \\ \text{subject to} \quad & 0 \leq x_1 \leq 1 \\ & \varepsilon x_{k-1} \leq x_k \leq 1 - \varepsilon x_{k-1} \quad \text{for } k = 2, \dots, n \\ & 0 \leq d + x_1 \quad \text{repeated } h_1 \text{ times} \\ & \varepsilon x_1 \leq d + x_2 \quad \text{repeated } h_2 \text{ times} \\ & \vdots \\ & \varepsilon x_{n-1} \leq d + x_n \quad \text{repeated } h_n \text{ times} \end{aligned}$$

To give a flavour of the main result, we first present Lemma 2.1 stating that, by adding $(d + 1)/(\varepsilon^{n-1}\delta)$ times the redundant inequality $\varepsilon x_{n-1} \leq d + x_n$ to the original KM formulation, the analytic centre χ^h can be pushed arbitrarily close to the vertex $v^{(n)} = (0, \dots, 0, 1)$.




Figure 3. The central path in the non-redundant Klee–Minty 2-cube.

To warranty, without loss of generality, that h is integer-valued, we assume that both $1/\varepsilon$ and $1/\delta$ are positive integers.

LEMMA 2.1 *Given $\delta \leq \varepsilon \leq 1/4$, d positive integer and $h = (0, \dots, 0, (d + 1)/(\varepsilon^{n-1}\delta))$, the analytic centre χ^h satisfies $|\chi^h - v^{[n]}|_\infty \leq \delta$.*

While Lemma 2.1 sets the starting point of the central path in the δ -neighbourhood of $v^{[n]}$, Proposition 2.2 states that, for a careful choice of d and h , the central path of the cube takes $2^n - 2$ turns before converging to the origin as it passes through the δ -neighbourhood of all the 2^n vertices of the Klee–Minty n -cube.

PROPOSITION 2.2 *Given $\varepsilon \leq 1/4$, $\delta < \varepsilon^{n-1}$, choose integer $d \geq n2^{n+1}$ and $h = 4nd/\delta$ ($(2^n - 1)/\varepsilon, \dots, (2^n - 2^{k-1})/\varepsilon^k, \dots, 2^{n-1}/\varepsilon^n$), then for each vertex v^S of the Klee–Minty n -cube, there is a point $\chi^h(v_n^S)$ of the central path satisfying $|\chi^h(v_n^S) - v^S|_\infty \leq \delta$.*

3. Proofs of Lemma 2.1 and Proposition 2.2

3.1 Proof of Lemma 2.1

The analytic centre $\chi^h = (\xi_1^h, \dots, \xi_n^h)$ of KM^h is the solution to the problem consisting of maximising the product of the slack variables

$$\begin{aligned} s_1 &= x_1 \\ s_k &= x_k - \varepsilon x_{k-1} && \text{for } k = 2, \dots, n \\ \bar{s}_1 &= 1 - x_1 \\ \bar{s}_k &= 1 - \varepsilon x_{k-1} - x_k && \text{for } k = 2, \dots, n \\ \tilde{s}_1 &= d + s_1 && \text{repeated } h_1 \text{ times} \\ &\vdots && \vdots \\ \tilde{s}_n &= d + s_n && \text{repeated } h_n \text{ times} \end{aligned}$$

Equivalently, χ^h is the solution of the following maximisation problem

$$\max_x \sum_{k=1}^n (\log s_k + \log \bar{s}_k + h_k \log \tilde{s}_k)$$

i.e. with the convention $x_0 = 0$

$$\max_x \sum_{k=1}^n (\log(x_k - \varepsilon x_{k-1}) + \log(1 - \varepsilon x_{k-1} - x_k) + h_k \log(d + x_k - \varepsilon x_{k-1}))$$

The optimality conditions (the gradient is equal to zero at optimality) for this concave maximisation problem give

$$\begin{aligned} \frac{1}{\sigma_k^h} - \frac{\varepsilon}{\sigma_{k+1}^h} - \frac{1}{\bar{\sigma}_k^h} - \frac{\varepsilon}{\bar{\sigma}_{k+1}^h} + \frac{h_k}{\tilde{\sigma}_k^h} - \frac{h_{k+1}\varepsilon}{\tilde{\sigma}_{k+1}^h} &= 0 \quad \text{for } k = 1, \dots, n - 1 \\ \frac{1}{\sigma_n^h} - \frac{1}{\bar{\sigma}_n^h} + \frac{h_n}{\tilde{\sigma}_n^h} &= 0 \\ \sigma_k^h > 0, \bar{\sigma}_k^h > 0, \tilde{\sigma}_k^h > 0 &\quad \text{for } k = 1, \dots, n, \end{aligned} \tag{1}$$

where

$$\begin{aligned} \sigma_1^h &= \xi_1^h \\ \sigma_k^h &= \xi_k^h - \varepsilon \xi_{k-1}^h \quad \text{for } k = 2, \dots, n \\ \bar{\sigma}_1^h &= 1 - \xi_1^h \\ \bar{\sigma}_k^h &= 1 - \varepsilon \xi_{k-1}^h - \xi_k^h \quad \text{for } k = 2, \dots, n \\ \tilde{\sigma}_k^h &= d + \sigma_k^h \quad \text{for } k = 1, \dots, n. \end{aligned}$$

The following lemma states that, for h_n large enough relatively to the other h_k values, the analytic centre χ^h is pushed to the neighbourhood of the vertex $v^{(n)} = (0, \dots, 0, 1)$.

LEMMA 3.1 *Given $\delta \leq \varepsilon \leq 1/4$ and h_1, \dots, h_{n-1} , we have*

$$|\chi^h - v^{(n)}|_\infty \leq \delta \quad \text{for } h_n \geq \frac{d+1}{\eta_n} \quad \text{where}$$

$$\frac{1}{\eta_n} = \max_{1 \leq k \leq n-1} \left\{ \frac{1}{2\varepsilon^{n-k}} \left(\frac{h_k}{d} + \frac{2}{\delta} \right) \right\}.$$

Proof The analytic centre $\chi^h = (\xi_1^h, \dots, \xi_n^h)$ is the solution of equation (1). Let us consider the n th equation of (1). As $\tilde{\sigma}_n^h \leq d+1$, we have $h_n \leq (d+1)/\tilde{\sigma}_n^h$. Thus, as $h_n \geq (d+1)/\eta_n$, we have $\tilde{\sigma}_n^h \leq \eta_n$, which implies $\tilde{\sigma}_n^h \leq \delta/2$. Let us then consider the $(n-1)$ th equation. We have

$$\frac{1}{\sigma_{n-1}^h} = \frac{1}{\bar{\sigma}_{n-1}^h} + \frac{\varepsilon}{\sigma_n^h} + \frac{\varepsilon}{\bar{\sigma}_n^h} - \frac{h_{n-1}}{\bar{\sigma}_{n-1}^h} + \frac{h_n \varepsilon}{\tilde{\sigma}_n^h} \geq \frac{\varepsilon}{\bar{\sigma}_n^h} - \frac{h_{n-1}}{\bar{\sigma}_{n-1}^h} + \frac{h_n \varepsilon}{\tilde{\sigma}_n^h}$$

As $\bar{\sigma}_n^h \leq \eta_n$, $\tilde{\sigma}_{n-1}^h \geq d$ and $\tilde{\sigma}_n^h \leq d+1$, this implies

$$\frac{1}{\sigma_{n-1}^h} \geq \frac{2\varepsilon}{\eta_n} - \frac{h_{n-1}}{d} \geq \frac{2}{\delta}$$

i.e. $\sigma_{n-1}^h \leq \delta/2$. The first $n-1$ equations of (1) can be rewritten as

$$\frac{1}{\sigma_k^h} + \frac{h_k}{\tilde{\sigma}_k^h} = \frac{1}{\bar{\sigma}_k^h} + \varepsilon \left(\frac{1}{\sigma_{k+1}^h} + \frac{h_{k+1}}{\tilde{\sigma}_{k+1}^h} \right) + \frac{\varepsilon}{\bar{\sigma}_{k+1}^h} \quad \text{for } k = 1, \dots, n-1.$$

For $k \leq n-2$, forward substitutions for the k th, $(k+1)$ th, \dots , $(n-1)$ th equations give

$$\frac{1}{\sigma_k^h} + \frac{h_k}{\tilde{\sigma}_k^h} = \frac{1}{\bar{\sigma}_k^h} + 2 \sum_{j=k+1}^{n-1} \frac{\varepsilon^{j-k}}{\bar{\sigma}_j^h} + \frac{\varepsilon^{n-k}}{\sigma_n^h} + \frac{h_n \varepsilon^{n-k}}{\tilde{\sigma}_n^h} + \frac{\varepsilon^{n-k}}{\bar{\sigma}_n^h}$$

which implies

$$\frac{1}{\sigma_k^h} \geq \frac{\varepsilon^{n-k}}{\bar{\sigma}_n^h} - \frac{h_k}{\tilde{\sigma}_k^h} + \frac{h_n \varepsilon^{n-k}}{\tilde{\sigma}_n^h}$$

As $\bar{\sigma}_n^h \leq \eta_n$, $\tilde{\sigma}_k^h \geq d$ and $\tilde{\sigma}_n^h \leq d+1$, this implies

$$\frac{1}{\sigma_k^h} \geq \frac{2\varepsilon^{n-k}}{\eta_n} - \frac{h_k}{d} \geq \frac{2}{\delta}$$

i.e. $\sigma_k^h \leq \delta/2$ for $k = 1, \dots, n-2$. Therefore, for $h_n \geq (d+1)/\eta_n$, we have $\xi_k^h \leq \delta$ for $k = 1, \dots, n-1$ and $1 - \xi_n^h \leq \delta$. ■

Lemma 2.1 is a direct corollary of Lemma 3.1 with $h = (0, \dots, 0, h_n)$.

3.2 Proof of Proposition 2.2

3.2.1 Preliminary Lemmas

LEMMA 3.2 Given $\delta \leq \varepsilon \leq 1/4$ and integer $d \geq n2^{n+1}$, then $h = 4nd/\delta((2^n - 1)/\varepsilon, \dots, (2^n - 2^{k-1})/\varepsilon^k, \dots, 2^{n-1}/\varepsilon^n)$ is a positive and integer solution of $Ah \geq b$, where $b = 4n/\delta(1, \dots, 1)$ and

$$A = \begin{pmatrix} \frac{-1}{d} & 0 & 0 & 0 & \dots & 0 & \frac{2\varepsilon^{n-1}}{d+1} \\ \frac{1}{d+1} & \frac{-\varepsilon}{d} & 0 & 0 & \dots & 0 & 0 \\ \frac{-1}{d} & \frac{2\varepsilon}{d+1} & \frac{-\varepsilon^2}{d} & 0 & \dots & 0 & 0 \\ \frac{-1}{d} & \frac{-\varepsilon}{d(d+1)} & \frac{2\varepsilon^2}{d+1} & \frac{-\varepsilon^3}{d} & \dots & 0 & 0 \\ \vdots & \vdots & \vdots & \vdots & \ddots & \vdots & \vdots \\ \frac{-1}{d} & \frac{-\varepsilon}{d(d+1)} & \frac{-\varepsilon^2}{d(d+1)} & \frac{-\varepsilon^3}{d(d+1)} & \dots & \frac{-\varepsilon^{n-2}}{d} & 0 \\ \frac{-1}{d} & \frac{-\varepsilon}{d(d+1)} & \frac{-\varepsilon^2}{d(d+1)} & \frac{-\varepsilon^3}{d(d+1)} & \dots & \frac{2\varepsilon^{n-2}}{d+1} & \frac{-\varepsilon^{n-1}}{d} \end{pmatrix}$$

Proof Multiplying both sides of $Ah \geq b$ by $\delta\varepsilon(d+1)/4n$, we have

$$d(1 - \varepsilon) \geq 2^n - 1 + \varepsilon$$

$$d(1 - \varepsilon) \geq 2^n - 2 + \varepsilon$$

$$d(1 - \varepsilon) \geq 2^{n+1} - 5 + \varepsilon$$

$$d(1 - \varepsilon) \geq 2^n - 2^k + \sum_{j=1}^{k-1} (2^n - 2^{j-1}) + \varepsilon \quad \text{for } k = 4, \dots, n,$$

which, as $d(1 - \varepsilon) \geq n2^n$, is implied by the obvious conditions

$$n2^n \geq 2^n - 1 + \varepsilon$$

$$n2^n \geq 2^n - 2 + \varepsilon$$

$$n2^n \geq 2^{n+1} - 5 + \varepsilon$$

$$n2^n \geq k2^n - 3 \cdot 2^{k-1} + 1 + \varepsilon \quad \text{for } k = 4, \dots, n.$$

■

To ease the notations, we define, for $k = 1, \dots, n - 1$

$$\ell_k = \frac{h_k}{d+1} - \frac{h_{k+1}\varepsilon}{d}$$

$$u_k = \frac{h_k}{d} - \frac{h_{k+1}\varepsilon}{d+1}$$

LEMMA 3.3 *The system $Ah \geq b$ is equivalent to*

$$\begin{aligned} \frac{2h_n \varepsilon^{n-1}}{d+1} - \frac{h_1}{d} &\geq \frac{4n}{\delta} \\ \ell_1 &\geq \frac{4n}{\delta} \\ \ell_k \varepsilon^{k-1} - \sum_{j=1}^{k-1} u_j \varepsilon^{j-1} &\geq \frac{4n}{\delta} \quad \text{for } k = 2, \dots, n-1. \end{aligned}$$

Proof The first two inequalities are direct reformulations of the first two inequalities of the system $Ah \geq b$. For $k = 2, \dots, n-1$, the inequality $\ell_k \varepsilon^{k-1} - \sum_{j=1}^{k-1} u_j \varepsilon^{j-1} \geq 4n/\delta$ can be rewritten as:

$$\frac{2h_k \varepsilon^{k-1}}{d+1} - \frac{h_{k+1} \varepsilon^k}{d} - \frac{h_1}{d} - \frac{1}{d(d+1)} \sum_{j=2}^{k-1} h_j \varepsilon^{j-1} \geq \frac{4n}{\delta}.$$

■

COROLLARY 3.4 *For h satisfying the last $n-1$ inequalities of $Ah \geq b$, we have*

$$\ell_k \varepsilon^{k-1} \geq \frac{2^{k+1}n}{\delta} \quad \text{for } k = 1, \dots, n-1.$$

Proof The proof is by induction on k . We have $\ell_1 \geq 4n/\delta$ and the result follows by using $u_k \geq \ell_k$ in Lemma 3.3. ■

COROLLARY 3.5 *Given $\delta \leq \varepsilon \leq 1/4$ and a positive integer h satisfying the last $n-1$ inequalities of $Ah \geq b$, we have*

$$|\chi^h - v^{[n]}|_\infty \leq \delta \quad \text{for} \quad \frac{2h_n \varepsilon^{n-1}}{d+1} \geq \frac{h_1}{d} + \frac{2}{\delta}$$

Proof Corollary 3.4 implies $\ell_k \geq 0$ for $k = 1, \dots, n-1$. Hence, in Lemma 3.1 we have $1/\eta_n = (h_1/d + 2/d)/(2\varepsilon^{n-1})$, which gives the result. ■

The central path of KM^h can be defined as the set of analytic centres $\chi^h(\alpha) = (x_1^h, \dots, x_{n-1}^h, \alpha)$ of the intersection of the hyperplane $H_\alpha: x_n = \alpha$ with the feasible region of KM^h where $0 \leq \alpha \leq \xi_n^h$, see [6]. These intersections are called the α -level sets and $\chi^h(\alpha)$ is the solution of the following system

$$\begin{aligned} \frac{1}{s_k^h} - \frac{\varepsilon}{s_{k+1}^h} - \frac{1}{\tilde{s}_k^h} - \frac{\varepsilon}{\tilde{s}_{k+1}^h} + \frac{h_k}{\tilde{s}_k^h} - \frac{h_{k+1}\varepsilon}{\tilde{s}_{k+1}^h} &= 0 \quad \text{for } k = 1, \dots, n-1 \\ s_k^h > 0, \tilde{s}_k^h > 0, \tilde{s}_{k+1}^h > 0 &\quad \text{for } k = 1, \dots, n-1 \end{aligned} \tag{2}$$

where

$$\begin{aligned}
 s_1^h &= x_1^h \\
 s_k^h &= x_k^h - \varepsilon x_{k-1}^h \quad \text{for } k = 2, \dots, n-1 \\
 s_n^h &= \alpha - \varepsilon x_{n-1}^h \\
 \bar{s}_1^h &= 1 - x_1^h \\
 \bar{s}_k^h &= 1 - \varepsilon x_{k-1}^h - x_k^h \quad \text{for } k = 2, \dots, n-1 \\
 \bar{s}_n^h &= 1 - \alpha - \varepsilon x_{n-1}^h \\
 \tilde{s}_k^h &= d + s_k^h \quad \text{for } k = 1, \dots, n.
 \end{aligned}$$

In the rest of the paper, we assume that $\delta \leq \varepsilon^{n-1}$ and that a positive integer h satisfying $Ah \geq b$ is given. Corollary 3.5 implies that $1 - \delta \leq \xi_n^h$ and therefore we can consider the α -level set for $\alpha \leq 1 - \varepsilon^{n-1}$ as it implies $\alpha \leq \xi_n^h$.

LEMMA 3.6 *Given $\varepsilon \leq 1/4$, $\delta < \varepsilon^{n-1}$, integer $d \geq n2^{n+1}$ and a positive integer h satisfying $Ah \geq b$; for $0 \leq \alpha \leq 1 - \varepsilon^{n-1}$ and $k \notin \{1, n\}$, if the k th coordinate x_k^h of the analytic centre $\chi^h(\alpha)$ satisfies*

$$x_k^h \in [\varepsilon^{k-1} - t_k \varepsilon^{k-1} \delta, 1 - \varepsilon^{k-1} + t_k \varepsilon^{k-1} \delta],$$

$$\text{where } \begin{cases} t_1 = 1 \\ t_2 = 1 - \frac{2}{4n-1} \\ t_{k+1} = t_k - \frac{1}{n} \quad \text{for } k = 2, \dots, n-1, \end{cases}$$

then, $x_{\hat{k}}^h \geq 1 - \varepsilon$ for some \hat{k} smaller than or equal to $k - 1$.

Proof Assume to the contrary that the statement is false, i.e. $x_{\hat{k}}^h < 1 - \varepsilon$ for $\hat{k} = 1, \dots, k - 1$. Considering the first equation of (2) and successively using $x_1^h < 1 - \varepsilon$, $\delta < \varepsilon$ and Lemma 3.3, we have

$$\frac{\varepsilon}{s_2^h} + \frac{\varepsilon}{\bar{s}_2^h} = \frac{1}{s_1^h} - \frac{1}{\bar{s}_1^h} + \frac{h_1}{\tilde{s}_1^h} - \frac{h_2 \varepsilon}{\bar{\tilde{s}}_2^h} \geq -\frac{1}{\varepsilon} + \frac{h_1}{d+1} - \frac{h_2 \varepsilon}{d} \geq -\frac{1}{\delta} + \frac{h_1}{d+1} - \frac{h_2 \varepsilon}{d} \geq \frac{4n-1}{\delta}$$

which implies either

$$x_2^h \leq \varepsilon x_1^h + \frac{2\varepsilon\delta}{4n-1} < \varepsilon(1 - \varepsilon) + \frac{2\varepsilon\delta}{4n-1} \leq \varepsilon(1 - \delta) + \frac{2\varepsilon\delta}{4n-1} = \varepsilon - t_2 \varepsilon \delta,$$

or

$$x_2^h \geq 1 - \varepsilon x_1^h - \frac{2\varepsilon\delta}{4n-1} > 1 - \varepsilon(1 - \varepsilon) - \frac{2\varepsilon\delta}{4n-1} > 1 - \varepsilon.$$

As $x_2^h < 1 - \varepsilon$, this implies $x_2^h < \varepsilon - t_2 \varepsilon \delta$. Similarly, considering the \hat{k} th equation of (2) for $\hat{k} = 2, \dots, k - 2$, we have $x_{\hat{k}}^h < \varepsilon^{\hat{k}-1} - t_{\hat{k}} \varepsilon^{\hat{k}-1} \delta$. Considering the $(k - 1)$ st equation and

successively using $x_{k-1}^h < \varepsilon^{k-1}$, $x_{k-2}^h < 1$ and Corollary 3.4, we have

$$\begin{aligned} \frac{\varepsilon}{s_k^h} + \frac{\varepsilon}{\bar{s}_k^h} &= \frac{1}{s_{k-1}^h} - \frac{1}{\bar{s}_{k-1}^h} + \frac{h_{k-1}}{\bar{s}_{k-1}^h} - \frac{h_k \varepsilon}{\bar{s}_k^h} \\ &\geq \frac{1}{\varepsilon^{k-2}} - \frac{1}{1 - \varepsilon^{k-2} - \varepsilon} + \frac{h_{k-1}}{d+1} - \frac{h_k \varepsilon}{d} \geq \ell_{k-1} \geq \frac{2^k n}{\varepsilon^{k-2} \delta} \end{aligned}$$

which implies either

$$x_k^h \leq \varepsilon x_{k-1}^h + \frac{\varepsilon^{k-1} \delta}{2^{k-1} n} < \varepsilon(\varepsilon^{k-2} - t_{k-1} \varepsilon^{k-2} \delta) + \frac{\varepsilon^{k-1} \delta}{2^{k-1} n} \leq \varepsilon^{k-1} - t_k \varepsilon^{k-1} \delta$$

or

$$x_k^h \geq 1 - \varepsilon x_{k-1}^h - \frac{\varepsilon^{k-1} \delta}{2^{k-1} n} > 1 - \varepsilon(\varepsilon^{k-2} - t_{k-1} \varepsilon^{k-2} \delta) - \frac{\varepsilon^{k-1} \delta}{2^{k-1} n} \geq 1 - \varepsilon^{k-1} + t_k \varepsilon^{k-1} \delta.$$

This is impossible as $x_k^h \in [\varepsilon^{k-1} - t_k \varepsilon^{k-1} \delta, 1 - \varepsilon^{k-1} + t_k \varepsilon^{k-1} \delta]$. ■

LEMMA 3.7 Given $\varepsilon \leq 1/4$, $\delta < \varepsilon^{n-1}$, $d \geq n2^{n+1}$, a positive integer h satisfying $Ah \geq b$ and t_1, \dots, t_n as specified in Lemma 3.6; for $0 \leq \alpha \leq 1 - \varepsilon^{n-1}$ and $k \notin \{1, n\}$, if the k th coordinate x_k^h of the analytic centre $\chi^h(\alpha)$ satisfies

$$x_k^h \in [t_k \varepsilon^{k-1} \delta, 1 - t_k \varepsilon^{k-1} \delta]$$

then $s_k^h \geq \varepsilon^{\hat{k}-1} \delta / (2^{\hat{k}} n)$ for some \hat{k} smaller than or equal to $k - 1$.

Proof Assume to the contrary that the statement is false, i.e. $s_{\hat{k}}^h < (1/2^{\hat{k}} n) \varepsilon^{\hat{k}-1} \delta$ for $\hat{k} = 1, \dots, k - 1$. This implies

$$x_k^h < \frac{\varepsilon^{\hat{k}-1} \delta}{4n} \sum_{j=1}^{\hat{k}-1} \frac{1}{2^{j-2}} \quad \text{for } \hat{k} = 1, \dots, k - 1$$

Considering the $(k - 1)$ st equation of (2) and using $\bar{s}_{k-1}^h = 1 - 2\varepsilon x_{k-2}^h - s_{k-1}^h$, we have

$$\begin{aligned} \frac{\varepsilon}{s_k^h} + \frac{\varepsilon}{\bar{s}_k^h} &= \frac{1}{s_{k-1}^h} - \frac{1}{\bar{s}_{k-1}^h} + \frac{h_{k-1}}{\bar{s}_{k-1}^h} - \frac{h_k \varepsilon}{\bar{s}_k^h} \\ &\geq \frac{2^{k-1} n}{\varepsilon^{k-2} \delta} - \frac{1}{1 - 2\varepsilon - 1/(2^{k-1} n) \varepsilon^{k-2} \delta} + \frac{h_{k-1}}{d+1} - \frac{h_k \varepsilon}{d} \geq \ell_{k-1} \end{aligned}$$

By Corollary 3.4, this implies

$$\frac{\varepsilon}{s_k^h} + \frac{\varepsilon}{\bar{s}_k^h} \geq \frac{2^k n}{\varepsilon^{k-2} \delta},$$

which further implies either, since $1/n \leq t_k$

$$\begin{aligned} x_k^h &\leq \varepsilon x_{k-1}^h + \frac{\varepsilon^{k-1} \delta}{2^{k-1} n} < \frac{\varepsilon^{k-1} \delta}{4n} \sum_{j=1}^{k-2} \frac{1}{2^{j-2}} + \frac{\varepsilon^{k-1} \delta}{2^{k-1} n} \\ &= \frac{\varepsilon^{k-1} \delta}{4n} \sum_{j=1}^{k-1} \frac{1}{2^{j-2}} \leq \frac{\varepsilon^{k-1} \delta}{n} \leq t_k \varepsilon^{k-1} \delta, \end{aligned}$$

or

$$\begin{aligned} x_k^h &\geq 1 - \varepsilon x_{k-1}^h - \frac{\varepsilon^{k-1}\delta}{2^{k-1}n} > 1 - \frac{\varepsilon^{k-1}\delta}{4n} \sum_{j=1}^{k-2} \frac{1}{2^{j-2}} - \frac{\varepsilon^{k-1}\delta}{2^{k-1}n} \\ &= 1 - \frac{\varepsilon^{k-1}\delta}{4n} \sum_{j=1}^{k-1} \frac{1}{2^{j-2}} \geq 1 - t_k \varepsilon^{k-1} \delta. \end{aligned}$$

This is impossible because $x_k^h \in [t_k \varepsilon^{k-1} \delta, 1 - t_k \varepsilon^{k-1} \delta]$. ■

3.2.2 Proof of Proposition 2.2 By analogy with the unit cube $[0, 1]^n$, we denote the vertices of the Klee–Minty cube \mathcal{C} using a subset S of $\{1, \dots, n\}$. For $S \subseteq \{1, \dots, n\}$, a vertex v^S of \mathcal{C} is defined by

$$\begin{aligned} v_1^S &= \begin{cases} 1, & \text{if } 1 \in S \\ 0, & \text{otherwise} \end{cases} \\ v_k^S &= \begin{cases} 1 - \varepsilon v_{k-1}^S, & \text{if } k \in S \\ \varepsilon v_{k-1}^S, & \text{otherwise} \end{cases} \quad k = 2, \dots, n. \end{aligned}$$

PROPOSITION 3.8 Given $\varepsilon \leq 1/4$, $\delta < \varepsilon^{n-1}$, $d \geq n2^{n+1}$ and a positive integer h satisfying $Ah \geq b$, for $k \neq n$, the $(k + 1)$ th and k th coordinates of the analytic centre $\chi^h(v_n^S)$ of the v_n^S -level set satisfy

$$|x_{k+1}^h - v_{k+1}^S| \leq t_{k+1} \varepsilon^k \delta \implies |x_k^h - v_k^S| \leq t_k \varepsilon^{k-1} \delta.$$

Proof Assume to the contrary that the statement is false, i.e. for at least one k smaller than or equal to $n - 1$, we have $|x_{k+1}^h - v_{k+1}^S| \leq t_{k+1} \varepsilon^k \delta$ and $|x_k^h - v_k^S| > t_k \varepsilon^{k-1} \delta$. We consider a case by case analysis.

Case 1 $v_k^S = 0$

The inequality $|x_k^h - v_k^S| > t_k \varepsilon^{k-1} \delta$ implies $x_k^h > t_k \varepsilon^{k-1} \delta$ and, as $\varepsilon x_k^h \leq x_{k+1}^h \leq 1 - \varepsilon x_k^h$, we have

$$t_k \varepsilon^k \delta < x_{k+1}^h < 1 - t_k \varepsilon^k \delta$$

As $t_{k+1} < t_k$, this implies $t_{k+1} \varepsilon^k \delta < x_{k+1}^h < 1 - t_{k+1} \varepsilon^k \delta$. This contradicts the inequality $|x_{k+1}^h - v_{k+1}^S| \leq t_{k+1} \varepsilon^k \delta$, where $v_{k+1}^S = 0$ or 1 because $v_k^S = 0$.

Case 2 $0 < v_k^S < 1$

The inequality $|x_k^h - v_k^S| > t_k \varepsilon^{k-1} \delta$ implies $x_k^h \in]0, v_k^S - t_k \varepsilon^{k-1} \delta[$ or $x_k^h \in]v_k^S + t_k \varepsilon^{k-1} \delta, 1[$. By $]a, b[$ we denote the open interval between a and b .

(1) $x_k^h \in]v_k^S + t_k \varepsilon^{k-1} \delta, 1[$

As $\varepsilon x_k^h \leq x_{k+1}^h \leq 1 - \varepsilon x_k^h$, we have $\varepsilon(v_k^S + t_k \varepsilon^{k-1} \delta) < x_{k+1}^h < 1 - \varepsilon(v_k^S + t_k \varepsilon^{k-1} \delta)$.

As $t_{k+1} < t_k$, this implies $\varepsilon v_k^S + t_{k+1} \varepsilon^k \delta < x_{k+1}^h < 1 - \varepsilon v_k^S - t_{k+1} \varepsilon^k \delta$. This contradicts the inequality $|x_{k+1}^h - v_{k+1}^S| \leq t_{k+1} \varepsilon^k \delta$, where $v_{k+1}^S = \varepsilon v_k^S$ or $1 - \varepsilon v_k^S$.

(2) $x_k^h \in]0, v_k^S - t_k \varepsilon^{k-1} \delta[$

(a) $x_k^h < \varepsilon^{k-1} - t_k \varepsilon^{k-1} \delta$

Considering the k th equation of (2) and successively using $x_k < \varepsilon^{k-1}$ and Corollary 3.4, we have

$$\begin{aligned} \frac{\varepsilon}{s_{k+1}^h} + \frac{\varepsilon}{\bar{s}_{k+1}^h} &= \frac{1}{s_k^h} - \frac{1}{\bar{s}_k^h} + \frac{h_k}{\bar{s}_k^h} - \frac{h_{k+1}\varepsilon}{\bar{s}_{k+1}^h} \\ &\geq \frac{1}{\varepsilon^{k-1}} - \frac{1}{1 - \varepsilon^{k-1} - \varepsilon} + \frac{h_k}{d+1} - \frac{h_{k+1}\varepsilon}{d} \geq \ell_k \geq \frac{2^{k+1}n}{\varepsilon^{k-1}\delta}, \end{aligned}$$

which implies either

$$x_{k+1}^h \leq \varepsilon x_k^h + \frac{1}{2^k n} \varepsilon^k \delta < \varepsilon(\varepsilon^{k-1} - t_k \varepsilon^{k-1} \delta) + \frac{1}{2^k n} \varepsilon^k \delta \leq \varepsilon^k - t_{k+1} \varepsilon^k \delta$$

or

$$x_{k+1}^h \geq 1 - \varepsilon x_k^h - \frac{\varepsilon^k \delta}{2^k n} > 1 - \varepsilon(\varepsilon^{k-1} - t_k \varepsilon^{k-1} \delta) - \frac{\varepsilon^k \delta}{2^k n} \geq 1 - \varepsilon^k + t_{k+1} \varepsilon^k \delta.$$

This contradicts the inequality $|x_{k+1}^h - v_{k+1}^S| \leq t_{k+1} \varepsilon^k \delta$, where $v_{k+1}^S \geq \varepsilon^k$ because $v_k^S > 0$.

(b) $\varepsilon^{k-1} - t_k \varepsilon^{k-1} \delta \leq x_k^h < v_k^S - t_k \varepsilon^{k-1} \delta$ (we have $k \neq 1$ as $0 < v_k^S < 1$)

By Lemma 3.6, there is a \hat{k} smaller than or equal to $k - 1$ such that $x_{\hat{k}}^h \geq 1 - \varepsilon$, which implies $s_{\hat{k}}^h \geq 1 - 2\varepsilon$. Considering the \hat{k} -th equation of (2) and using $s_{\hat{k}}^h \geq 1 - 2\varepsilon$, we have

$$\begin{aligned} \frac{\varepsilon^{\hat{k}}}{s_{\hat{k}+1}^h} &= \frac{\varepsilon^{\hat{k}-1}}{s_{\hat{k}}^h} - \frac{\varepsilon^{\hat{k}-1}}{\bar{s}_{\hat{k}}^h} - \frac{\varepsilon^{\hat{k}}}{\bar{s}_{\hat{k}+1}^h} + \frac{h_{\hat{k}}\varepsilon^{\hat{k}-1}}{\bar{s}_{\hat{k}}^h} - \frac{h_{\hat{k}+1}\varepsilon^{\hat{k}}}{\bar{s}_{\hat{k}+1}^h} \\ &\leq \frac{\varepsilon^{\hat{k}-1}}{1 - 2\varepsilon} - \frac{\varepsilon^{\hat{k}-1}}{\varepsilon} - \frac{\varepsilon^{\hat{k}}}{\bar{s}_{\hat{k}+1}^h} + \frac{h_{\hat{k}}\varepsilon^{\hat{k}-1}}{d} - \frac{h_{\hat{k}+1}\varepsilon^{\hat{k}}}{d+1} \end{aligned}$$

which implies

$$\frac{\varepsilon^{\hat{k}}}{s_{\hat{k}+1}^h} \leq \frac{h_{\hat{k}}\varepsilon^{\hat{k}-1}}{d} - \frac{h_{\hat{k}+1}\varepsilon^{\hat{k}}}{d+1} = u_{\hat{k}}\varepsilon^{\hat{k}-1}$$

The previous inequality, which corresponds to the case $i = \hat{k}$, can be generalised to

$$\frac{\varepsilon^i}{s_{i+1}^h} \leq \sum_{j=\hat{k}}^i u_j \varepsilon^{j-1} \quad \text{for } i = \hat{k}, \dots, k - 1$$

which clearly holds for $k = \hat{k} + 1$ and, for $k > \hat{k} + 1$, is obtained by multiplying the i th equation of (2) by ε^{i-1} for $i = \hat{k} + 1, \dots, k - 1$ and using successively a similar argument.

Noticing that we could have initially permute $s_{\hat{k}+1}^h$ and $\bar{s}_{\hat{k}+1}^h$, gives

$$\frac{\varepsilon^{k-1}}{\bar{s}_k^h} \leq \sum_{j=\hat{k}}^{k-1} u_j \varepsilon^{j-1}$$

Together with the k th equation of (2), this implies

$$\frac{\varepsilon^k}{s_{k+1}^h} + \frac{\varepsilon^k}{\bar{s}_{k+1}^h} = \frac{\varepsilon^{k-1}}{s_k^h} - \frac{\varepsilon^{k-1}}{\bar{s}_k^h} + \frac{h_k \varepsilon^{k-1}}{\bar{s}_k^h} - \frac{h_{k+1} \varepsilon^k}{\bar{s}_{k+1}^h} \geq \frac{h_k \varepsilon^{k-1}}{d+1} - \frac{h_{k+1} \varepsilon^k}{d} - \sum_{j=\hat{k}}^{k-1} u_j \varepsilon^{j-1}$$

i.e.

$$\frac{\varepsilon^k}{s_{k+1}^h} + \frac{\varepsilon^k}{\bar{s}_{k+1}^h} \geq \ell_k \varepsilon^{k-1} - \sum_{j=\hat{k}}^{k-1} u_j \varepsilon^{j-1}$$

Using Lemma 3.3, Corollary 3.4 and $u_k \geq \ell_k$, this implies

$$\frac{\varepsilon^k}{s_{k+1}^h} + \frac{\varepsilon^k}{\bar{s}_{k+1}^h} \geq \frac{4n}{\delta} + \sum_{j=1}^{\hat{k}-1} u_j \varepsilon^{j-1} \geq \frac{2^{\hat{k}+1} n}{\delta}$$

which implies either

$$x_{k+1}^h \leq \varepsilon x_k^h + \frac{\varepsilon^k \delta}{2^{\hat{k}n}} < \varepsilon(v_k^S - t_k \varepsilon^{k-1} \delta) + \frac{\varepsilon^k \delta}{2^{\hat{k}n}} \leq \varepsilon v_k^S - t_{k+1} \varepsilon^k \delta,$$

or

$$x_{k+1}^h \geq 1 - \varepsilon x_k^h - \frac{\varepsilon^k \delta}{2^{\hat{k}n}} > 1 - \varepsilon(v_k^S - t_k \varepsilon^{k-1} \delta) - \frac{\varepsilon^k \delta}{2^{\hat{k}n}} \geq 1 - \varepsilon v_k^S + t_{k+1} \varepsilon^k \delta.$$

This contradicts the inequality $|x_{k+1}^h - v_{k+1}^S| \leq t_{k+1} \varepsilon^k \delta$, where $v_{k+1}^S = \varepsilon v_k^S$ or $1 - \varepsilon v_k^S$.

Case 3 $v_k^S = 1$

The inequality $|x_k^h - v_k^S| > t_k \varepsilon^{k-1} \delta$ implies $x_k^h < 1 - t_k \varepsilon^{k-1} \delta$.

(1) $x_k^h < t_k \varepsilon^{k-1} \delta$

Considering the k th equation of (2) and successively using $x_k^h < t_k \varepsilon^{k-1} \delta$, $t_k \varepsilon^{k-1} \delta \leq \varepsilon$ and Corollary 3.4, we have

$$\begin{aligned} \frac{\varepsilon}{s_{k+1}^h} + \frac{\varepsilon}{\bar{s}_{k+1}^h} &= \frac{1}{s_k^h} - \frac{1}{\bar{s}_k^h} + \frac{h_k}{\bar{s}_k^h} - \frac{h_{k+1} \varepsilon}{\bar{s}_{k+1}^h} \\ &\geq \frac{1}{t_k \varepsilon^{k-1} \delta} - \frac{1}{1 - t_k \varepsilon^{k-1} \delta - \varepsilon} + \frac{h_k}{d+1} - \frac{h_{k+1} \varepsilon}{d} \geq \ell_k \geq \frac{2^{k+1} n}{\varepsilon^{k-1} \delta} \end{aligned}$$

which implies either

$$x_{k+1}^h \leq \varepsilon x_k^h + \frac{\varepsilon^k \delta}{2^{\hat{k}n}} < t_k \varepsilon^k \delta + \frac{\varepsilon^k \delta}{2^{\hat{k}n}} = \left(t_k + \frac{1}{2^{\hat{k}n}} \right) \varepsilon^k \delta$$

or

$$x_{k+1}^h \geq 1 - \varepsilon x_k^h - \frac{\varepsilon^k \delta}{2^{\hat{k}n}} > 1 - t_k \varepsilon^k \delta - \frac{\varepsilon^k \delta}{2^{\hat{k}n}} = 1 - \left(t_k + \frac{1}{2^{\hat{k}n}} \right) \varepsilon^k \delta$$

This contradicts the inequality $|x_{k+1}^h - v_{k+1}^S| \leq t_{k+1} \varepsilon^k \delta$, where $v_{k+1}^S = \varepsilon$ or $1 - \varepsilon$ because $v_k^S = 1$.

(2) $t_k \varepsilon^{k-1} \delta \leq x_k^h < 1 - t_k \varepsilon^{k-1} \delta$

(a) $k = 1$

From the first equation of (2), we have

$$\frac{\varepsilon}{s_2^h} + \frac{\varepsilon}{\bar{s}_2^h} = \frac{1}{s_1^h} - \frac{1}{\bar{s}_1^h} + \frac{h_1}{\bar{s}_1^h} - \frac{h_2 \varepsilon}{\bar{s}_2^h} \geq -\frac{1}{\delta} + \ell_1 \geq \frac{4n-1}{\delta}$$

which implies either

$$x_2^h \leq \varepsilon x_1^h + \frac{2\varepsilon\delta}{4n-1} < \varepsilon(1-\delta) + \frac{2\varepsilon\delta}{4n-1} \leq \varepsilon - t_2 \varepsilon \delta$$

or

$$x_2^h \geq 1 - \varepsilon x_1^h - \frac{2\varepsilon\delta}{4n-1} > 1 - \varepsilon(1-\delta) - \frac{2\varepsilon\delta}{4n-1} \geq 1 - \varepsilon + t_2 \varepsilon \delta$$

This contradicts $|x_2^h - v_2^S| \leq t_2 \varepsilon \delta$ where $v_2^S = \varepsilon$ or $1 - \varepsilon$ as $v_1^S = 1$.

(b) $k \neq 1$

By Lemma 3.7, there is a \hat{k} smaller than or equal to $k - 1$ such that $s_{\hat{k}}^h \geq \varepsilon^{\hat{k}-1} \delta / (2^{\hat{k}} n)$. Considering the \hat{k} th equation of (2), we have

$$\begin{aligned} \frac{\varepsilon^{\hat{k}}}{s_{\hat{k}+1}^h} &= \frac{\varepsilon^{\hat{k}-1}}{s_{\hat{k}}^h} - \frac{\varepsilon^{\hat{k}-1}}{\bar{s}_{\hat{k}}^h} - \frac{\varepsilon^{\hat{k}}}{\bar{s}_{\hat{k}+1}^h} + \frac{h_{\hat{k}} \varepsilon^{\hat{k}-1}}{\bar{s}_{\hat{k}}^h} - \frac{h_{\hat{k}+1} \varepsilon^{\hat{k}}}{\bar{s}_{\hat{k}+1}^h} \\ &\leq \frac{2^{\hat{k}} n}{\delta} + \frac{h_{\hat{k}} \varepsilon^{\hat{k}-1}}{d} - \frac{h_{\hat{k}+1} \varepsilon^{\hat{k}}}{d+1} = u_{\hat{k}} \varepsilon^{\hat{k}-1} + \frac{2^{\hat{k}} n}{\delta} \end{aligned}$$

This inequality, which corresponds to the case $i = \hat{k}$, can be generalised to

$$\frac{\varepsilon^i}{s_{i+1}^h} \leq \sum_{j=\hat{k}}^i u_j \varepsilon^{j-1} + \frac{2^{\hat{k}} n}{\delta} \quad \text{for } i = \hat{k}, \dots, k-1,$$

which clearly holds for $k = \hat{k} + 1$ and, for $k > \hat{k} + 1$, is obtained by multiplying the i th equation of (2) by ε^{i-1} for $i = \hat{k} + 1, \dots, k - 1$ and using successively a similar argument. Noticing that we could have initially permute $s_{\hat{k}+1}^h$ and $\bar{s}_{\hat{k}+1}^h$, gives

$$\frac{\varepsilon^{k-1}}{\bar{s}_k^h} \leq \sum_{j=\hat{k}}^{k-1} u_j \varepsilon^{j-1} + \frac{2^{\hat{k}} n}{\delta}$$

Together with the k th equation of (2), this implies

$$\begin{aligned} \frac{\varepsilon^k}{s_{k+1}^h} + \frac{\varepsilon^k}{\bar{s}_{k+1}^h} &= \frac{\varepsilon^{k-1}}{s_k^h} - \frac{\varepsilon^{k-1}}{\bar{s}_k^h} + \frac{h_k \varepsilon^{k-1}}{\bar{s}_k^h} - \frac{h_{k+1} \varepsilon^k}{\bar{s}_{k+1}^h} \\ &\geq -\sum_{i=\hat{k}}^{k-1} u_i \varepsilon^{i-1} - \frac{2^{\hat{k}} n}{\delta} + \frac{h_k \varepsilon^{k-1}}{d+1} - \frac{h_{k+1} \varepsilon^k}{d}. \end{aligned}$$

Using Lemma 3.3, Corollary 3.4 and $u_k \geq \ell_k$, the previous inequality gives

$$\begin{aligned} \frac{\varepsilon^k}{s_{k+1}^h} + \frac{\varepsilon^k}{\bar{s}_{k+1}^h} &\geq \ell_k \varepsilon^{k-1} - \sum_{i=\hat{k}}^{k-1} u_i \varepsilon^{i-1} - \frac{2^{\hat{k}} n}{\delta} \\ &\geq \frac{4n}{\delta} + \sum_{i=1}^{\hat{k}-1} u_i \varepsilon^{i-1} - \frac{2^{\hat{k}} n}{\delta} \geq \frac{2^{\hat{k}+1} n}{\delta} - \frac{2^{\hat{k}} n}{\delta} = \frac{2^{\hat{k}} n}{\delta} \end{aligned}$$

which implies either

$$x_{k+1}^h \leq \varepsilon x_k^h + \frac{\varepsilon^k \delta}{2^{\hat{k}-1} n} < \varepsilon(1 - t_k \varepsilon^{k-1} \delta) + \frac{\varepsilon^k \delta}{2^{\hat{k}-1} n} \leq \varepsilon - t_{k+1} \varepsilon^k \delta$$

or

$$x_{k+1}^h \geq 1 - \varepsilon x_k^h - \frac{\varepsilon^k \delta}{2^{\hat{k}-1} n} > 1 - \varepsilon(1 - t_k \varepsilon^{k-1} \delta) - \frac{\varepsilon^k \delta}{2^{\hat{k}-1} n} \geq 1 - \varepsilon + t_{k+1} \varepsilon^k \delta$$

This contradicts $|x_{k+1}^h - v_{k+1}^S| \leq t_{k+1} \varepsilon^k \delta$ where $v_{k+1}^S = \varepsilon$ or $1 - \varepsilon$ as $v_k^S = 1$. ■

Proposition 2.2 is a direct corollary of Proposition 1 as, for $S \neq \emptyset$ and $S \neq \{n\}$, we have $|x_n^h - v_n^S| = 0$; implying $|x_k^h - v_k^S| \leq t_k \varepsilon^{k-1} \delta$ for $k = 1, \dots, n - 1$. In other words, $|\chi^h(v_n^S) - v^S|_\infty \leq \delta$. Furthermore, by Corollary 3.5 we have $|\chi^h - v^{\{n\}}|_\infty \leq \delta$, and the central path converges to the origin v^\emptyset .

4. Remarks and future work

1. We showed that, without changing the geometry of the feasible set of KM, the central path can be forced to visit arbitrarily small neighbourhoods of all the vertices of the Klee–Minty n -cube by carefully adding redundant constraints.
2. This result highlights that, although the central path is a smooth analytical curve in the interior of the set of feasible solutions, it might be severely distracted by redundant constraints. In particular, exponentially many redundant constraints interplaying with the geometry of the problem, may force the central path to take exponentially many and arbitrarily sharp turns.
3. Our example leads to an $\Omega(2^n)$ lower bound for the number of iterations needed for central path-following interior point methods. The theoretical iteration-complexity upper bound $O(\sqrt{NL}) = O(2^{9n} n^4)$ as, for this example, the number of constraints $N = O(2^{6n} n^2)$ and the bit length of the input data $L = O(2^{6n} n^3)$. Therefore, the $\Omega(2^n)$ lower bound yields an $\Omega(\sqrt[6]{N/\ln^2 N})$ iteration-complexity lower bound. Using a different analysis, Todd and Ye [9] gave an $\Omega(\sqrt[3]{N})$ iteration-complexity lower bound between two updates of the barrier function. In a subsequent paper, Deza *et al.* [10] essentially closed the gap between the lower and upper bounds.
4. State-of-the-art preprocessing tools in modern linear optimisation software would eliminate the added redundant inequalities. Therefore, interior point methods based codes would solve the preprocessed KM^h efficiently, just as commercial simplex codes do solve the KM in only one pivot. A challenging task would be to design a variant of KM^h that cannot be easily simplified by known preprocessing heuristics.

Acknowledgements

Research supported by the NSERC Discovery grant number 48923 and a MITACS grant for the last three authors, by the Canada Research Chair program for the first and last authors and by the NSERC Discovery grant number 311969 for the first author.

References

- [1] Dantzig, G.B., 1951 Maximization of a linear function of variables subject to linear inequalities. In: T.C. Koopmans (Ed.) *Activity Analysis of Production and Allocation* (New York: John Wiley), pp. 339–347.
- [2] Klee, V. and Minty, G.J., 1972, How good is the simplex algorithm? In: O. Shisha (Ed.) *Inequalities III*, (Boston: Academic Press), pp. 159–175.
- [3] Terlaky, T. and Zhang, S. 1993, Pivot rules for linear programming – a survey. *Annals of Operations Research*, **46**, 203–233.
- [4] Khachiyan, L.G., 1979, A polynomial algorithm in linear programming. *Soviet Mathematics Doklady*, **20**, 191–194.
- [5] Karmarkar, N.K., 1984, A new polynomial-time algorithm for linear programming. *Combinatorica* **4**, 373–395.
- [6] Roos, C., Terlaky, T. and Vial, J-Ph., 1997, *Theory and Algorithms for Linear Optimization: An Interior Point Approach*. Wiley-Interscience Series in Discrete Mathematics and Optimization (New York: John Wiley).
- [7] Ye, Y., 1979, *Interior-Point Algorithms: Theory and Analysis*. Wiley-Interscience Series in Discrete Mathematics and Optimization (New York: John Wiley).
- [8] Wright, S.J., 1997, *Primal-Dual Interior-Point Methods* (Philadelphia: SIAM Publications).
- [9] Todd, M. and Ye, Y., 1996, A lower bound on the number of iterations of long-step and polynomial interior-point linear programming algorithms. *Annals of Operations Research*, **62**, 233–252.
- [10] Deza, A. Nematollahi, E. and Terlaky, T., 2004, How good are interior point methods? Klee–Minty cubes tighten iteration–complexity bounds. AdvOL-Report 2004/20, McMaster University.
- [11] Megiddo, N., 1986, Pathways to the optimal set in linear programming. In: N. Megiddo (Ed.) *Progress in Mathematical Programming: Interior-Point and Related Methods* (New York: Springer-Verlag), pp. 131–158; also in: Proceedings of the 7th Mathematical Programming Symposium of Japan, 1986, Nagoya, Japan, pp. 1–35.
- [12] Sonnevend, G. 1985, An “analytical centre” for polyhedrons and new classes of global algorithms for linear (smooth, convex) programming. In: A. Prékopa, J. Szelezsán and B. Strazicky (Eds) *System Modelling and Optimization: Proceedings of the 12th IFIP-Conference*, Budapest 1985. Also Lecture Notes in Control and Information Sciences 84, 1986 (New York: Springer), pp. 866–876.